

## Background material

---

1

## Relations

---

- A relation over a set  $S$  is a set  $R \subseteq S \times S$ 
  - We write  $a R b$  for  $(a,b) \in R$
- A relation  $R$  is:
  - reflexive iff  $\forall a \in S . a R a$
  - transitive iff  $\forall a \in S, b \in S, c \in S . a R b \wedge b R c \Rightarrow a R c$
  - symmetric iff  $\forall a, b \in S . a R b \Rightarrow b R a$
  - anti-symmetric iff  $\forall a, b, \in S . a R b \Rightarrow \neg(b R a)$

2

## Relations

---

- A relation over a set  $S$  is a set  $R \subseteq S \times S$ 
  - We write  $a R b$  for  $(a,b) \in R$
- A relation  $R$  is:
  - reflexive iff  $\forall a \in S . a R a$
  - transitive iff  $\forall a \in S, b \in S, c \in S . a R b \wedge b R c \Rightarrow a R c$
  - symmetric iff  $\forall a, b \in S . a R b \Rightarrow b R a$
  - anti-symmetric iff  $\forall a, b, \in S . a R b \Rightarrow \neg(b R a)$
  - $\forall a, b, \in S . a R b \wedge b R a \Rightarrow a = b$

3

## Partial orders

---

- An equivalence class is a relation that is:
- A partial order is a relation that is:

4

## Partial orders

---

- An equivalence class is a relation that is:
  - reflexive, transitive, symmetric
- A partial order is a relation that is:
  - reflexive, transitive, anti-symmetric
- A partially ordered set (a poset) is a pair  $(S, \leq)$  of a set  $S$  and a partial order  $\leq$  over the set
- Examples of posets:  $(2^S, \subseteq)$ ,  $(\mathbb{Z}, \leq)$ ,  $(\mathbb{Z}, \text{divides})$

5

## Lub and glb

---

- Given a poset  $(S, \leq)$ , and two elements  $a \in S$  and  $b \in S$ , then the:
  - least upper bound (lub) is an element  $c$  such that  $a \leq c, b \leq c$ , and  $\forall d \in S . (a \leq d \wedge b \leq d) \Rightarrow c \leq d$
  - greatest lower bound (glb) is an element  $c$  such that  $c \leq a, c \leq b$ , and  $\forall d \in S . (d \leq a \wedge d \leq b) \Rightarrow d \leq c$
- Does a lub and glb always exists?
  - A. Yes (in this case justify your answer)
  - B. No (in this case come up with example where no lub or glb exists)

6

## Lub and glb

- Given a poset  $(S, \leq)$ , and two elements  $a \in S$  and  $b \in S$ , then the:
  - least upper bound (lub) is an element  $c$  such that  $a \leq c$ ,  $b \leq c$ , and  $\forall d \in S. (a \leq d \wedge b \leq d) \Rightarrow c \leq d$
  - greatest lower bound (glb) is an element  $c$  such that  $c \leq a$ ,  $c \leq b$ , and  $\forall d \in S. (d \leq a \wedge d \leq b) \Rightarrow d \leq c$
- lub and glb don't always exist:



7


## Lattices

- A lattice is a tuple  $(S, \sqsubseteq, \perp, \top, \sqcup, \sqcap)$  such that:
  - $(S, \sqsubseteq)$  is a poset
  - $\forall a \in S. \perp \sqsubseteq a$
  - $\forall a \in S. a \sqsubseteq \top$
  - Every two elements from  $S$  have a lub and a glb
  - $\sqcup$  is the least upper bound operator, called a join
  - $\sqcap$  is the greatest lower bound operator, called a meet

8

## Examples of lattices


- Powerset lattice



9

## Examples of lattices


- Powerset lattice



10

## Examples of lattices


- Booleans expressions



11

## Examples of lattices

- Booleans expressions



12

## Examples of lattices

- Booleans expressions

two ways

T

⋮

F

13

## Examples of lattices

- Booleans expressions

two ways

T

⋮ 16 branches...

F

$(2^n)$   

2	0	1	0
1	0	0	0
1	1	0	0

14